

34 - STABILIZING MUSCLES OF THE LUMBAR SPINE STRENGTHENING IN THE TREATMENT OF LOW BACK PAIN

ANDRESSA MAYRA DOS SANTOS¹

MARINÊZ BOEING RUARO²

ANDERSOM RICARDO FRÉZ³

JOÃO AFONSO RUARO⁴

1. Physiotherapist/FAA; 2. Physiotherapist, Postural and Manual Therapy Specialist/CESUMAR;

3. Physiotherapist, Master in Health and Interdisciplinary Rehabilitation/UNICAMP;

4. Physiotherapist, Doctorating in Health Sciences/UFRN. UFRN

Clinic School of Physiotherapy, Faculty Anglo-Americano/FAA, Foz do Iguacu, Parana, Brazil.

Professor.joaoruaro@facisa.ufrn.br

INTRODUCTION

Low back pain is a common disorder, debilitating and costly financially^{1, 2}. Epidemiological studies estimate that the prevalence of low back pain in the general population is between 50 and 80%³. It is known that the etiology of localized pains column is multifactorial and may result from mechanical causes, degenerative factors or by psychosomatic⁴ disorders, ⁵, and one of the changes associated with low back pain is muscle imbalance.

As an important risk factor for low back pain is muscle weakness of the muscles of the trunk and abdomen⁶⁻⁸. The strengthening of these muscles appears to be a viable alternative for the management of dor^{9, 10}, including associated control nutritional¹¹, the fisioterapia¹², the use of ultrassom¹³ suíça¹⁴ and ball.

The two muscles most involved in these cases appear to be the lumbar multifidus (LM) and transverse abdominis (TA)¹⁵. Pesquisas¹⁶⁻¹⁸ show that muscle dysfunction is a ML after the first episode of low back pain, the muscle mechanoreceptors deconditions and commits, they could not recover spontaneously, even with the reduction of pain. In healthy subjects the TA is contracted before the movements of the extremities to protect the column, while in individuals suffering from back pain before this contraction movements failed, triggering an inefficient muscular stabilization of the spine, thus requiring a treatment específico^{19, 20}.

Figueiredo et al.²¹ cite three studies which were using needle electromyography and ultrasound real-time results show that the TA can be contracted in advance and in priority to other abdominal muscles during the depression of the abdominal wall, but despite the effectiveness of the equipment considered "gold standard", they have some limitations.

Still, according to the authors, due to the difficulty of using these laboratory devices in rehabilitation centers, it is essential to develop clinical techniques for evaluation and training of TA, one possibility being the use of Biofeedback Unit pressure (UBP), as the stabilizer. The UBP is a device developed by physiotherapists, which aims to quantify changes in pressure in an inelastic bag, sandwiched between the abdomen and the stretcher, during contraction of the TA, a correct contraction of the muscle in question resulted in a change in pressure recorded manômetro²².

The TA should be trained separately as one of the major muscles in the affected low back pain, losing its function tônica²³ because its contraction through specific exercise significantly reduces the laxity of the sacroiliac joint, hence the use of such contractions may be useful for the treatment of lombalgia²⁴.

In addition, this training provides the individual a stabilizer improves the automatic stabilizers synergism of other muscles, being rich in proprioceptive stimulation, giving it an improved performance of their atividades²⁵.

Based on the dysfunctions of the stabilizing muscles of the spine, this research was to evaluate the effect of an exercise protocol in the treatment of low back pain, emphasizing the strengthening of the TA muscle in order to provide the patient with greater body awareness and efficiency of muscle contraction, by monitoring with the UBP.

MATERIALS AND METHODS

This research was characterized as an experimental study of randomized clinical trial. The study was conducted on the premises of the School of Physical Therapy Clinic of the Anglo-American School of Foz do Iguacu, in the period from September to October 2009.

The sample was initially composed by 21 subjects, aged between 20 and 60 years, of both genders, with a clinical diagnosis of low back pain. However, three subjects were excluded from the sample because they had herniated discs. Thus, the remaining 18 subjects were randomly divided into two groups: treatment group (TG) and control group (CG).

The GT was subjected to an exercise protocol based on the method of segmental stabilization through proprioceptive exercises focusing on TA muscle, while the CG did not receive the intervention, was observed only with the intention of checking the natural evolution of the pathology.

Three subjects dropped out of the GT treatment for personal reasons. Thus, the GT at the end was composed of six of nine subjects and the GC.

Both groups underwent an initial assessment including: 1) identification and history, 2) completing the questionnaire McGill²⁶ pain, which is composed of 78 descriptors that evaluate the characteristics of pain and its intensity, 3) completing the questionnaire of Roland -morris²⁷, which consists of 24 items on the interference of back pain in activities of daily living and life skills, and 4) test transverse abdominal muscle activation, through the use of the stabilizer unit, brand Chattanooga²², inflating the manometer to 70 mmHg and requesting the contraction of the TA.

Stabilizer for the use of the patient was positioned prone, with the device in the region of the TA, held after inspiration and expiration associated with contraction of isolated muscle, a contraction of the TA correct results in reduced blood pressure greater than or equal to 4 mmHg²¹.

After application of the exercise protocol, both groups were reevaluated. One researcher was responsible for the history and application of questionnaires, both the beginning and end of the protocol, while another researcher performed the measurements of TA muscle activation.

Research volunteers were instructed to practice the exercises below, aimed at stabilizing muscle proprioception of the lumbar spine and postural positioning.

The protocol was applied for 25 sessions, divided into three phases: I) preparation (1 to 25 of treatment) and ii) two stages of specific training: cognitive and associative (6 th to 15 th service), and III) stage autonomous and functional training (16 ° to 25 ° care). The proposed therapy was applied three times a week, with an average of 40 minutes each, always observing the importance of palpation and verbal command.

The following describes each phase of the exercises:

Phase I: a) heating and mobility: the patient was asked to perform scales anterior-posterior and lateral-lateral hip on the Swiss ball, repeating 10 times every movement, and rotational motion on the Swiss ball (10 times to the right and 10 to the left) (adapted from Craig28), b) manual therapy (technical gear): patient prone, rolling apply T10 to the sacral region for 2 minutes, bilaterally to the spinous processes (adapted from Guirro and Guirro29), c) diaphragmatic breathing (Sarmento30), repeating 5 times and controlled via palpation and verbal command.

Phase II (cognitive stage): a) patient in the supine position, asked if the contraction of the rectus abdominis, TA later and eventually muscle contraction of the perineum, 5 repetitions. b) held a three-simultaneous contraction (rectum and perineum TA) contraction for 5 seconds and relax for 10 seconds, 10 repetitions.

Phase II (associative stage): three-simultaneous contraction of muscle training and relaxation for 10 seconds 5 seconds, in two different positions: sitting patient and extent of the lower limbs cinnamon 1 kg (5 repetitions for each member) and patient standing and half-squat (about 30 degrees of knee flexion and trunk leaning against the wall, 10 repetitions).

Phase III (autonomous stage and functional training): a) tri-simultaneous contraction of the muscles in training, 20 seconds of contraction and relaxation 10 b) motion to sit / stand, then full squats leaning on backrest with 5 repetitions each c) mini-circuit: walk 10 meters with obstacles, climbing ramp and fall down the stairs (stair-ramp), take 1 kg object on the floor, load it and put it back on the floor, filling the bladder (balloon) and secure it in place above the shoulders (2 repetitions).

The study was approved by the Ethics Committee of Faculty and Research Assis Gurgacz, Opinion no. 251/2009. All signed an informed consent.

The results were obtained by statistical analysis with the Student t test and Wilcoxon test, using Graphpad Instat software version 3.05. We calculated the mean values, standard deviation and values obtained with the subtraction of the values collected in the pre-and post-implementation of the McGill questionnaire, Roland-Morris and contraction of the test performed by device stabilizer.

RESULTS

Table I presents the data of the study. You can see the values obtained with medium and (difference) between the GT and GC, respectively, before the application of forms of assessment used.

Table I: Comparison of mean values and standard deviation obtained through the evaluation forms used.

	Treated group		Control group		Δ between GT and GC		
	Pre	Post	Pre	Pre	Post	Pre	P
McGill	48,7±4,9	39,2±10,4	40,3±9,4	41,4±7,9	9,5±10,9	-1,1±7,9	0,0599
Roland-Morris	13,7±4,3	5,2±3,1	8,9±5,1	9,6±5,1	8,5±3,9	-0,7±1,1	<0,001*
Stabilizaer	85,0±3,2	65,3±2,5	80,1±5,0	80,8±5,4	19,7±5,1	-0,7±1,2	<0,001*

*:Statistically significant

After statistical analysis of comparisons values (difference) between the GT and GC, referring to the McGill questionnaire data, the p value was 0.0599, considered not significant.

However, when comparing the values obtained between the GT and GC with the Roland-Morris questionnaire, and test the transversus abdominis muscle contraction, using the stabilizer, we found values of p <0.001 for both considered extremely significant.

DISCUSSION

There is evidence that activities related to motor control, implemented as treatment programs and exercícios12 including physiotherapy, manual therapy and reeducação31 or containing only manual32 therapy are more effective than drug treatment in cases of chronic low back pain.

Similar events were observed in systematic reviews of Van Tulder et al.33 and Ferreira et al.34, who found that exercise therapy is more effective than usual medical approach in treating chronic back pain, in order to improve pain, the function and quality of life, including how to prevent episodes of recurrence.

These findings with the results of this work, especially with regard to the function back, after all the Roland Morris questionnaire represents an important tool for evaluating patients with low back pain, especially in studies clínicos27.

The literature review carried out by Gouveia and Gouveia20 clarified and strengthened the role of the TA muscle as a stabilizer back, its relationship with low back pain and the importance of specific training to improve their function.

Santos and Freitas25 also include other muscles responsible for stability, diaphragm, multifidus and pelvic floor. In the work of Fozatti et al.35, the authors observed that the balance between lumbar lordosis, the TA muscle, diaphragmatic motion and the axis of rotation of the coxofemoral joints, can be considered as a basic principle of treating stress incontinence by postural reeducation. Thus, the importance of the TA muscle exercises focused not only have benefits for low back pain.

Commenting on the acceptance of treatment, Wilson et al.36 reported that despite the contraction of these muscle groups increase the stiffness of the trunk with minimal load to the lumbar spine, the technique is not always easy for all patients, a fact which was not observed in this study. The subjects had no difficulty when performing the exercises.

In the review of France et al.15 The authors report that the exercises of segmental stabilization in low back pain have an important role in preventing the recurrence of low back pain, because these exercises restore the protective function of the deep muscles, and are specific to avoid overloading damaged structures. This fact could not be observed in this study, because the monitoring of the subjects occurred only during the treatments.

Figueiredo et al.21 evaluated the intra-examiner and between the use of the stabilizer during contraction of the TA muscle and concluded that the use of this tool should be encouraged in clinical practice because of its relevance, simplicity and safety reasons. A similar conclusion also found in the study of Costa et al.37, which describe the use of tests and palpation of the stabilizer for the analysis of TA muscle activation is reliable, even emphasizing that these methods of analysis should be encouraged because of its relevance, simplicity and safety reasons.

Following the line of the studies cited during this study sought to keep all procedures in a standardized way to avoid possible errors in the tests, including the placement of volunteers, calibration of the UBP and the movements that could influence the results, as respiratory movements, for example. In addition, the examiner underwent prior training to minimize possible errors.

In assessing pain, McGill26 questionnaire is widely used, including research linking the use of exercises on the ground or to the aquatic environment in the treatment program of low back pain, such as Norris and Matthews38, Waller et al. 39

and Hough et al.40. However, despite this variable in evolution with reduction of the GT score of the questionnaire in relation to GC after applying the proposed protocol of exercises, there was no statistical significance.

CONCLUSION

The proposed protocol of exercises to strengthen the stabilizing muscles of the lumbar spine resulted in improvement of lumbar function, improved muscle activation and TA did not significantly interfere in pain.

REFERENCES

- PATEL, S
1.; BROWN, S.; FRIEDE, T.; GRIFFITHS, F.; LORD, J.; NGUNJIRI, A.; THISTLETHWAITE, J.; TYSALL, C.; WOOLVINE, M.; UNDERWOOD, M. Study protocol: Improving patient choice in treating low back pain (IMPACT - LBP): a randomised controlled trial of a decision support package for use in physical therapy. **BMC Musculoskeletal Disorders**, n. 12, v. 52, p. 1-7, 2011.
- MASTERS, S
2.; LIND, R. Musculoskeletal pain - presentations to general practice. **Australian Family Physicians** v. 39, n. 6, p. 425-428, 2010.
- 3.HELFFENSTEIN JUNIOR M, GOLDENFUM MA, SIENA C. Occupational low back pain. **Revista da Associação Médica Brasileira**, v. 56, n. 5, p. 583-589, 2010.
- LAERUM, E
4.; BROX, J.I.; WERNER, E.L. Low back pain-still a clinical challenge. **Tidsskr Nor Laegeforen**, v. 130, n. 22, p. 2248-2251, 2010.
- SIKIRU, L
5.; HANIFA, S. Prevalence and risk factors of low back pain among nurses in a typical Nigerian hospital. **African Health Sciences**, v. 10, n. 1, p. 26-30, 2010.
- GRUTHER, W
6.; WICK, F.; PAUL, B.; LEITNER, C.; POSCH, M.; MATZNER, M.; CREVENNA, R.; EBENBICHLER, G. Diagnostic accuracy and reliability of muscle strength and endurance measurements in patients with chronic low back pain. **Journal of Rehabilitation Medicine**, v. 41, n. 8, p. 613-619, 2009.
- BAYRAMOĞLU, M
7.; AKMAN, M.N.; KILINÇ, S.; CETIN, N.; YAVUZ, N.; OZKER, R. Isokinetic measurement of trunk muscle strength in women with chronic low-back pain. **American Journal of Physcial Medicine and Rehabilitation**, v. 80, n. 9, p. 650-655, 2001.
- OLIVIER, N
8.; LEPRETRE, A.; CABY, I.; DUPUIS, M.A.; PRIEUR, F. Does exercise therapy for chronic lower-back pain require daily isokinetic reinforcement of the trunk muscles? **Annales de Readaptation et de Medecine Physique**, v. 51, n. 4, p. 284-291, 2005.
- BARR, K.P
9.; GRIGGS, M.; CADBY, T. Lumbar stabilization: a review of core concepts and current literature, part 2. **American Journal of Physcial Medicine and Rehabilitation**, v. 86, n. 1, p. 71-80, 2007.
- HEBERT, J.J
10.; KOPPENHAVER, S.L.; MAGEL, J.S.; FRITZ, J.M. The relationship of transversus abdominis and lumbar multifidus activation and prognostic factors for clinical success with a stabilization exercise program: a cross-sectional study. **Archives of Physical Medicine and Rehabilitation**, v. 91, n. 1, p. 78-85, 2010.
- ROFFEY, D.M
11.; ASHDOWN, L.C.; DORNAN, H.D.; CREECH, M.J.; DAGENAIS, S.; DENT, R.M.; WAI, E.K. Pilot evaluation of a multidisciplinary, medically supervised, nonsurgical weight loss program on the severity of low back pain in obese adults. **The Spine Journal**, v. 11, n. 3, p. 197-204, 2011.
- KO, J.K
12.. Comparing the effects of drug therapy, physical therapy, and exercise on pain, disability, and depression in patients with chronic low back pain. **Taehan Kanho Hakhoe Chi**, v. 37, n. 5, p. 645-654, 2007.
- GOREN, A
13.; YILDIZ, N.; TOPUZ, O.; FINDIKOGLU, G.; ARDIC, F. Efficacy of exercise and ultrasound in patients with lumbar spinal stenosis: a prospective randomized controlled trial. **Clinical Rehabilitation**, v. 24, n. 7, p. 623-631, 2010.
- ESCAMILLA, R.F
14.; LEWIS, C.; BELL, D.; BRAMBLET, G.; DAFFRON, J.; LAMBERT, S.; PECSON, A.; IMAMURA, R.; PAULO, L.; ANDREWS, J.R. Core muscle activation during Swiss ball and traditional abdominal exercises. **Journal of Orthopaedic and Sports Physical Therapy**, v. 40, n. 5, p. 265-276, 2010.
- 15.FRANÇA, F.J.R.; BURKE, T.N.; CLARET, D.C.; MARQUES, A.P. Estabilização segmentar da coluna lombar nas lombalgias: uma revisão bibliográfica e um programa de exercícios. **Fisioterapia e Pesquisa**, v. 15, n. 2, p. 200-206, 2008.
- 16.HIDES, J.A.; RICHARDSON, C.A.; JULL, G.A. Multifidus muscle recovery is not automatic following resolution of acute first-episode low back pain. **Spine**, v. 21, n. 23, p. 2763-2769, 1996.
- 17.HIDES, J.A.; STOKES, M.J.; SAIDE, M.; JULL, G.A.; COOPER, D.H. Evidence of lumbar multifidus muscle wasting ipsilateral to symptoms in patients with acute/subacute low back pain. **Spine**, n. 19, p. 165-172, 1994.
- 18.STOKES, M.; YOUNG, A. Investigations of quadriceps inhibition: implications for clinical practice. **Physiotherapy**, n. 70, p. 425-428, 1984.
- 19.SALMELLA, L.F.T.; SAKAMOTO, A.C.L.; SIQUEIRA, F.B. Mecanismos de estabilização da coluna lombar: uma revisão de literatura. **Fisioterapia em Movimento**, v. 17, n. 4, p. 51-58, 2004.
- 20.GOUVEIA, K.M.C.; GOUVEIA, E.C. O músculo transverso abdominal e sua função de estabilização da coluna lombar. **Fisioterapia em Movimento**, v. 21, n. 3, p. 45-50, 2008.
- 21.FIGUEIREDO, M.K.; CHAVES JÚNIOR, I.P.; FIGUEIREDO, V.G.C.; COSTA, L.O.P.; COSTA, L.C.M. Estudo da confiabilidade intra e entre-examinadores da unidade de biofeedback pressórico na medida da contração do músculo transverso abdominal. **Revista Brasileira de Ciência e Movimento**, v. 13, n. 4, p. 93-100, 2005.
- 22.ENCORE MEDICAL. Stabilizer Pressure Biofeedback. Operating Instructions. Brisbane: Chattanooga Pacific, 2002.
- 23.HODGES, P. Is there a role for transverses abdominis in lumbo-pelvic stability? **Manual Therapy**, v. 4, n. 2, p. 74-

86, 1999.

24. RICHARDSON, C.A.; SNIJDERS, C.J.; HIDES, J.A.; DAMEN, L.; PAS, M.S.; STORM, J. The relationship between the transversus abdominis muscles, sacroiliac joint mechanics, and low back pain. **Spine**, v. 27, n. 4, p. 399-405, 2002.

25. SANTOS, J.P.M.; FREITAS, G.F.P. Métodos de treinamento da estabilização central. **Semina**, v. 31, n. 1, p. 93-101, 2010.

26. PIMENTA, C.A.M.; TEIXEIRA, M.J. Questionário de dor McGill: proposta de adaptação para a língua portuguesa. **Revista da Escola de Enfermagem da USP**, v. 30, n. 3, p. 473-483, 1996.

27. NUSBAUM, L.; NATOUR, J.; FERRAZ, M.B.; GOLDENBERG, J. Translation, adaptation and validation of the Roland-Morris questionnaire - Brazil Roland-Morris. **Brazilian Journal of Medical and Biological Research**, v. 34, n. 2, p. 203-210, 2001.

28. CRAIG, C. **Pilates com a bola**. 2 ed. São Paulo: Phorte, 2005.

29. GUIRRO, E.C.O.; GUIRRO, R.R.J. **Fisioterapia dermatofuncional: fundamentos, recursos, patologias**. 3 ed. Barueri: Manole, 2004.

30. SARMENTO, G.J.V. **Fisioterapia respiratória no paciente crítico: rotinas clínicas**. 2 ed. Barueri: Manole, 2007.

31. MOSELEY, L. Combined physiotherapy and education is efficacious for chronic low-back pain. **The Australian Journal of Physiotherapy**, n. 48, p. 297-301, 2002.

32. NIEMISTO, L.; LAHTINEN-SUOPANKI, T.; RISSANEN, P.; LINDGREN, K.A.; SARNA, S.; HURRI, H. A randomized trial of combined manipulation, stabilizing exercises, and physician consultation compared to physician consultation alone for chronic low-back pain. **Spine**, n. 28, p. 2185-2191, 2003.

33. VAN TULDER, M.; FURLAN, A.; BOMBARDIER, C.; BOUTER, L. Update method guidelines for systematic reviews in the Cochrane Collaboration Back Review Group. **Spine**, n. 28, p. 1290-1299, 2003.

34. FERREIRA, M.C.; PENIDO, H.; AUN, A.; FERREIRA, P.; FERREIRA, M.L.; OLIVEIRA, V.C. Eficácia dos exercícios de controle motor na dor lombopélvica: uma revisão sistemática. **Fisioterapia e Pesquisa**, v. 16, n. 4, p. 374-379, 2009.

35. FOZZATTI, M.C.M.; PALMA, P.; HERRMANN, V.; DAMBROS, M. Impacto da reeducação postural global no tratamento da incontinência urinária de esforço feminina. **Revista da Associação Médica Brasileira**, v. 54, n. 1, p. 17-22, 2008.

36. WILSON, J.D.; DOUGHERTY, C.P.; IRELAND, M.L.; DAVIS, I.M. Core stability and relationship to lower extremity function and injury. **Journal of the American Academy of Orthopaedic Surgeons**, v. 13, n. 5, p. 316-325, 2005.

37. COSTA, L.O.P.; COSTA, L.C.M.; CANÇADO, R.L.; OLIVEIRA, W.M.; FERREIRA, P.H. Confiabilidade do teste palpatório e da unidade de biofeedback pressórico na ativação do músculo transverso abdominal em indivíduos normais. **Acta Fisiátrica**, v. 11, n. 3, p. 101-105, 2004.

NORRIS, C

38. MATTHEWS, M. The role of an integrated back stability program in patients with chronic low back pain. **Complementary Therapies in Clinical Practice**, v. 14, n. 4, p. 255-263, 2008

WALLER, B

39. LAMBECK, J.; DALY, D. Therapeutic aquatic exercise in the treatment of low back pain: a systematic review. **Clinical Rehabilitation**, v. 23, n. 1, p. 3-14, 2009.

HOUGHÉ

40. STEPHENSON R, SWIFT L. A comparison of manual therapy and active rehabilitation in the treatment of non specific low back pain with particular reference to a patient's Linton & Hallden psychological screening score: a pilot study. **BMC Musculoskeletal Disorders**, v. 1, n. 8, p. 106, 2007.

Autor correspondente: JOÃO AFONSO RUARO

Faculdade de Ciências da Saúde / Universidade Federal do Rio Grande do Norte – FACISA/UFRN.

Rua Trairi, s/n – Santa Cruz/RN CEP: 59200-000.

Telefone: (84)9650-1000

joaoruaro@facisa.ufrn.br

STABILIZING MUSCLES OF THE LUMBAR SPINE STRENGTHENING IN THE TREATMENT OF LOW BACK

PAIN

ABSTRACT

Background: The high prevalence in the population and the significant negative social and economic impact generated by the cases of disability justify the importance of the study of low back pain. Objective: Evaluate the effect of an exercise protocol to strengthen the stabilizing muscles of the lumbar spine in the treatment of low back pain. Methods: The proposed protocol was applied in 15 subjects randomly assigned to two groups, treated (TG) and a control group (CG), aged between 20 and 60. Methods of assessment questionnaires were used to McGill and Roland-Morris and stabilizer pressure biofeedback unit. The TG was submitted to an exercise protocol, based on the segmental stabilization, which emphasizes the strengthening of the transversus abdominis muscle. The CG did not receive the intervention. Results and conclusion: The results were that the TG presented significant improvement of lumbar function, the transverse abdominal muscle activation (both $p < 0.0001$) and clinical improvement of pain, but not statistically significant ($p = 0.0599$).

KEYWORDS: low back pain; exercise therapy, physical therapy.

RENFORCEMENT DES MUSCLES STABILISATEURS DE LA COLONNE LOMBAIRE LOMBAIRE DANS LE TRAITEMENT DE LOMBALGIES

SOMMAIRE

Introduction: La prévalence élevée dans la population et l'impact négatif significatif socioéconomiques engendrés par les cas de handicap ne justifie l'importance de l'étude de la douleur dorsale. Objectif: évaluer l'effet d'un protocole d'exercice pour renforcer les muscles stabilisateurs de la colonne lombaire dans le traitement de la lombalgie. Méthodologie: Le projet de protocole a été appliqué dans 15 sujets répartis aléatoirement en deux groupes, l'un traité (GT) et un groupe contrôle (GC), âgés entre 20 et 60. Comme les méthodes de questionnaires d'évaluation ont été utilisés de McGill et de Roland-Morris et stabilisateur de biofeedback pression. Le GT a été soumis à un protocole d'exercice basé sur la méthode de stabilisation segmentaire, en insistant sur le renforcement musculaire abdominale transversale. Le CG n'a pas bénéficié de l'intervention. Résultats et

conclusion: Les résultats ont montré que la GT a été une amélioration statistiquement significative de la fonction de retour, l'activation du muscle transverse de l'abdomen (les deux $p < 0,0001$) et l'amélioration clinique de la douleur, mais non statistiquement significative ($p = 0,0599$).

MOTS-CLÉS: douleurs lombaires, une rééducation, la physiothérapie.

FORTALECIMIENTO DE LOS MUSCULOS ESTABILIZADORES DE LA COLUMNA LUMBAR EN EL TRATAMIENTO DEL DOLOR DE ESPALDA

RESUMEN

Introducción: La alta prevalencia en la población y los importantes efectos socioeconómicos negativos generados por los casos de discapacidad justifican la importancia del estudio del dolor de espalda. Objetivo: Evaluar el efecto de un protocolo de ejercicios para fortalecer los músculos estabilizadores de la columna lumbar en el tratamiento del dolor de espalda baja.

Metodología: El proyecto de protocolo se aplicó en 15 pacientes divididos aleatoriamente en dos grupos, uno tratado (GT) y un grupo control (GC), con edades comprendidas entre 20 y 60. Como métodos de cuestionarios de evaluación se utilizaron McGill y Roland Morris y unidad de estabilización de la presión de biofeedback. El GT fue sometido a un protocolo de ejercicio basado en el método de estabilización segmentaria, haciendo hincapié en el fortalecimiento transversal de los músculos abdominales. La CG no recibió la intervención. Resultados y conclusiones: Los resultados mostraron que el GT fue una mejoría estadísticamente significativa en función de la espalda, la activación del músculo transverso del abdomen (ambos $p < 0,0001$) y la mejoría clínica del dolor, pero no estadísticamente significativo ($p = 0,0599$).

PALABRAS-CLAVE: dolor de espalda, terapia de ejercicios de fisioterapia.

FORTALECIMIENTO DOS MÚSCULOS ESTABILIZADORES DA COLUNA LOMBAR NO TRATAMENTO DA LOMBALGIA

RESUMO

Introdução: A alta prevalência na população e o expressivo impacto socioeconômico negativo gerado pelos casos de incapacidade justificam a importância do estudo das dores lombares. Objetivo: Avaliar o efeito de um protocolo de exercícios de fortalecimento dos músculos estabilizadores da coluna lombar no tratamento da lombalgia. Metodologia: O protocolo proposto foi aplicado em 15 sujeitos divididos aleatoriamente em dois grupos, um tratado (GT) e um controle (GC), com faixa etária entre 20 e 60 anos. Como métodos de avaliação foram utilizados os questionários de McGill e Roland-Morris e a unidade de biofeedback pressórico estabilizador. O GT foi submetido a um protocolo de exercícios, baseados no método de estabilização segmentar, enfatizando o fortalecimento do músculo transverso abdominal. O GC não recebeu a intervenção. Resultados e conclusão: Os resultados obtidos foram que o GT apresentou melhora estatisticamente significativa da função lombar, da ativação do músculo transverso abdominal (ambos com $p < 0,0001$) e melhora clínica da dor, porém não estatisticamente significativa ($p = 0,0599$).

PALAVRAS-CHAVE: dor lombar; terapia por exercício, fisioterapia.