

79 - COMBINED EFFECT OF TRAINING IN HYPERTENSIVE INDIVIDUAL WITH OVERWEIGHT, POST-ABLATION OF WOLFF-PARKINSON-WHITE SYNDROME, ON FACTORS THAT TRIGGER ATRIAL FIBRILLATION

FÁBIO PROCÓPIO SOARES BENTO
 RAFAEL SILVA GALDINO MATHIAS
 GLEISSON ALISSON PEREIRA DE BRITO
 Graduate Program Latu Sensu Gama Filho
 University in Obesity and Weight Loss
 Belo Horizonte-MG-Brazil
 fabiomapex@yahoo.com.br

INTRODUCTION

Quality of life can be considered as resulting from a set of parameters individual, social, cultural and environmental conditions that characterize the living being human, a community or a nation (Nahas, 2001 cited in Dantas, 2005).

According to Wilmore and Costill (2001), millions of people die of hunger every year most of the world, starting from many Americans are dying by the consumption excessive food. The authors also report that they are currently spending billions of dollars with overeating, which consequently leads to an expenditure of billions of dollars in weight loss methods.

Obesity is a multifactorial disease that may result from several factors genetic, physiological, environmental and psychological, which can provide the accumulation excessive energy in the form of fat in the body. (Yadav et al. 2001 apud Al Barbosa 2008). This abnormal accumulation of fat represents health risk and establishes a relationship with heart disease, such as hypertension.

"Atrial fibrillation (AF) is a supraventricular arrhythmia that occurs in a complete disorganized atrial electrical activity, causing the atria lose their ability contraction, not generating atrial systole" (Zimmerman, 2009). The Wolff-Parkinson-White (WPW) syndrome is characterized by a short PR interval, the appearance of a wave and Delta by the lengthening of the QRS, according to the ACLS (2007).

There are now studies showing the effectiveness of exercise in control hypertension and in improving the quality of life of individuals with AF. As Rique say, Meirelles and Soares (2002), in his work on Nutrition and Exercise in Control of Cardiovascular Diseases, regular exercise has a therapeutic role important in the control of systolic hypertension (SH) and, although the mechanisms responsible are not yet fully defined.

As cited by Generoso and Navarro (2011), the American College of Medicine Sports, reports the importance of physical activity associated with health and has promoted the results of the training against resistance in several studies to pathologies including hypertension.

This study aimed to examine the effects of an exercise program aerobic and muscular endurance on cardiovascular parameters in an individual with overweight, hypertension, post-ablation of Wolff-Parkinson-White about the factors that trigger atrial fibrillation.

MATERIALS AND METHODS

This case study was conducted on the premises of the Academy Fields - SESC in the city of Teófilo Otoni, Minas Gerais. We studied the individual VPS, sex male, 57 years, weight 80 kg, BMI 26.8, hypertensive patients with Wolff-Parkinson-White (WPW).

Periodic physical evaluations were performed during the training period that was of 3 months. We carried out measurement of heart rate and blood pressure measurements before, during and after training sessions.

Work with evaluation consisted of anthropometric measurements (weight, height, circumference) and skinfold thickness. Being weighed only swim trunks and barefoot, in a scale scientific Welmy brand, the measure of height was measured in the very balance using a rigid metal rod that reaches a height of 2 meters intervals in cm, so with the individual positioned back. The folds were measured with adipometer scientist at Sanny, with verification of 9 skinfolds, measured 3 times, and the median of each withdrawal and a perimeter with a metal strip on a scale Sanny millimeter of 2 meters in length.

The data were analyzed statistically using t test Student, using a significance level of $p < 0.05$.

MEDICAL HISTORY

The individual VPS was diagnosed with Wolff-Parkinson-White conduction pathway with anomaly located in the posterior left with the heart in 2005, asymptomatic, primary characteristic of the syndrome by the year 2005 when the symptoms of disease vigorously expressed. At first received medical advice to move away from physical activities that the picture was until it was fully understood by ablation surgery.

In 2006 came the first VPS ablation surgery, followed by improved frame. However, in 2010 there was recurrence, with installation of similar figures being performed a new ablation procedure with cauterization and complete interruption of the road anomalous WPW.

VPS during surgery led to a very easy one episode of atrial fibrillation that maintained, and it was necessary to carry out procedures cardiovenção 4 (CVE) during the entire procedure.

The patient also makes use of the following drugs: Pantocal® - pantoprazole sodium, Hyperium® 1mg - rilmenidine 1mg, Symbicort® - formoterol fumarate dihydrate / budesonide and Atacand HCT® - candesartan cilexetil + hydrochlorothiazide.

HISTORY OF PHYSICAL ACTIVITY

VPS was marathon athlete in his youth from 1975 to 1991 and football player amateur since I was 18 years old. He was sedentary since 2005, when the request was Cardiologist to interrupt the activities due to the diagnosis of WPW.

EXPERIMENTAL DESIGN

PSV made three weekly workouts at the gym. The training started with aerobic program in the initial weeks with the aim of cardiorespiratory adaptation

Arriving at the gym the individual received the Polar heart monitor FS2c, had their blood pressure measured by a digital Sphygmomanometer Onida brand. In was then subjected to a stretching session and forwarded to the mat, where walked for 15 minutes. At the end of time had their blood pressure and heart rate checked again, then follow horizontally to the stationary bike, which performed 15 minutes of exercise. At the end of this period, was again measured blood pressure and heart rate. After

the training, there was a second session stretching.

After the adaptation phase lasting three weeks, began training weathered, with a muscular endurance program (RML), alternated by follow-up, with the first objective of strengthening tendons, ligaments and joints, to better quality and resistance in the evolution of training exercises with 8, 4 lower limb (LL) and 4 for upper limbs (ULs). This training is for maintain aerobic work being done in the form of circuit without rest.

RESULTS AND DISCUSSION

As the weeks following training, was visible improvement in the provision in training, heart rate has not climbed as fast, and he succeeded after a pause for one minute to make the low frequency faster than occurred before. Even still taking antihypertensive medications, in a few days isolated, the individual came to the gym with high BP, which associate with causes external.

It was noted an improvement in heart rate, which increased the intensity of wake of 5.0 km to 6.0 km, the frequency that stayed in an average of 105 bpm in 6.0km was at 91 bpm. After the time of 15 minutes on the treadmill, the frequency trophy rose to 100 bpm, not to exceed this level during the 15 minutes thereafter, which indicates an improvement in cardiovascular fitness after four weeks of training.

With the progression of training intensity VPS has also evolved. This did not affect their BP that remained controlled during training and later to it. Even with the cargo growth, the volume of training was the same and he offered no complaints fatigue, or feeling bad because of the physical effort. The hypotension caused by exercise the end of the training sessions at all affect the physiological condition of the same.

"The post-exercise hypotensive effect is associated with physiological changes, such as reflection modulator mechanism of clay, by hyperemia control of muscle contractions and suppression of sympathetic activity." (Generoso and Navarro, 2011).




Figure 2: (Photo by electrocardiogram VPS after the second ablation of 28/05/2010, where the QRS interval is normal, with normal PR)

Table I: Tracking the Evolution of Circles.

Measures	Neck	Shoulder	Pectoral	Waist	Abdomen	Hips	Arm	Forearm	Thigh	Calf
05/12/2010	38,6 cm	106,7 cm	100,2 cm	96,5 cm	96,2 cm	97 cm	31,4cm	26,8 cm	53,9 cm	36,6 cm
06/02/2011	38,6 cm	106,3 cm	101,3 cm	93,6 cm	94,6 cm	96 cm	31,1 cm	27,0 cm	54,3 cm	37,3 cm
26/03/2011	37,0 cm	105,0 cm	102 cm	90,4 cm	90,2 cm	95 cm	32,0 cm	28,1 cm	56,3 cm	38,0 cm

Table II: BMI, Height and Weight

Measurement	Date	BMI	Height	Weight
01	05/12/2010	26,1	1,75	80
02	06/02/2011	24,4	1,75	74,6
03	26/03/2011	23,3	1,75	71,3

Table III: Tracking the Evolution of the skin folds

Measurement	Triceps	Subscapular	Biceps	Chest	Axillary	Abdomen	Suprailiac	Thigh	Calf
05/12/2010	17,8	28,4	5,0	17,1	18,9	21,1	16,3	18,1	4,8
06/02/2011	16,5	27,8	4,5	16,5	18,2	20,2	15,5	16,7	4,2
26/03/2011	15,7	26,2	4,1	15,9	17,4	18,2	14,8	15,9	4,0




Figure 03.A. Average systolic blood pressure (SBP) and B. diastolic blood pressure (DBP); the first week of training (white bars) and in the last week of training (Striped bars). Data are presented as mean ± SEM. * P <0.05.




Figure 04. A. Average mean arterial pressure (MAP) and B. heart rate (HR) in first week of training (white bars) and last week of training (bars stripes). Data are presented as mean \pm SEM. * $P < 0.05$.

Tables I, II and III respectively show the evolution of the individual studied in relation to the circumference, height, weight and BMI and skinfold thickness. Where was the presenting weight reduction, reduction measures and possible increased fat-free muscle mass in some measurements of skinfolds.

Last week, SBP, MAP and HR (Fig. 03 and 04) were significantly smaller during training compared to the first week. The resting HR in Last week was also lower when compared to the first week of CF. The WFP is the pressure in a complete cardiac cycle, and is calculated as $(MAP = DAP + 1/3PP)$, where PP is the difference between SBP and DBP.

Physical activity is an important resource in the prevention of atherosclerotic disease and effective adjuvant treatment, through cardiac rehabilitation. (Ghorayeb and COLS, 1998)

In addition to aerobic exercise, resistance training activities have increased their importance in cardiac rehabilitation (Rique, Soares and Meirelles, 2002).

Wilmore and Costill (2001), argue that obesity may contribute to a greater development of certain diseases, and weight reduction usually prescribed as an integral part of treatment.

The reduction of overweight and intra-abdominal adiposity and improved insulin resistance exercise may be aided by other mechanisms of action of this on hypertension according to Rique, Soares and Meirelles, (2002).

CONCLUSION

The combined training with the individual VPS, which underwent a procedure ablation to block an anomalous pathway of WPW, and later presented a framework AF, was able to control BP and HR. The individual did not show significant increases BP during training or after it, thus controlling hypertension. A reduction in weight with change in the classification of overweight for a weight within the normal range, going from 26.1 to 23.3 BMI. These results show reduction or control factors associated with the propensity for AF, which are old, overweight, obesity, hypertension and sedentary lifestyle. Thus, we conclude that the need for physical exercise is eminent to regulate these factors and provide an improvement in living conditions for the individual suffering from atrial fibrillation, resulting in better quality of life and longevity.

REFERENCES

- ACLS, Advanced Cardiac Life Support / Barbara Aehlert, [scientific review Ana Paula Quilici et al. Translation of Alexander MaceriMida et al]. - Rio de Janeiro: Elsevier, 2007. 4 th edition.
- BARBOSA, Jose Bonifacio. Metabolic Syndrome Clinic of Cardiology in SaoLuis - Maranhao. Thesis - Postgraduate Program - Undergraduate Health Maternal and Child Health. Federal University of Maranhão. 2008
- DANTAS, HM Estélio Thinking the Body and Movement. 2nd ed. Rio de Janeiro: Shape, 2005.
- D'AVILA, André, SAAD, Eduardo. Catheter Ablation Syndrome Wolff-Parkinson-White. Center for Atrial Fibrillation. Arrhythmia Service and Cardiac Pacing. Hospital Pro- Heart Rate. Ergometry Department, Cardiology and Cardiac Rehabilitation of Sports SOCERJ, DERCAD / RJ. Available at: < <http://www.dercad.org.br/boletim/ano6num26/ablacao.asp> > Accessed: 01/07/2011.
- DE MATOS, L.D.N.; PASTORE, C.A.; SAMESIMA, N., FRANCO, Changes in FGDM Resting ECG with the Physical Training. IN: NEGRÃO, EC, Barreto, ACP Cardiology Exercise: Athlete to heart disease. 3rd Ed Barueri, SP: Manole, 2010.
- FISHER, Naomi D.L.; WILLIAMS, Gordon H. hypertensive vascular disease. Harrison Internal Medicine / editors Dennis L. Kasper ... [Et al.]. - 16. ed. - Rio de Janeiro: McGraw-Hill Inter-Brazil Ltda., 2006.
- FLATO, U.A.P., D'EVA, F.C., FLATO, E.M.S., Vomero, J.S., FERREIRA, D.M.; Guimarães, H.P.; LOPES, R.D. New Evidence on Antithrombotic Atrial Fibrillation 2010. Journal of Clinical Medicine.
- GENEROSO, Alexsandro Fernandes; NAVARRO, Antônio Coppi. Resistance training and promote aerobic regulation in blood pressure levels in a sedentary individual and hypertensive. Journal of Exercise Physiology, V10, No. 2, April / June 2011.
- GHORAYEB, N, BAPTISTA, CA, DIOGUARDI, GS, CHIGI, A., STEPHAN, D., DAHER, D., SMITH, P. Exercise in Patients with Atrial Fibrillation. Journal of the Society Cardiology of São Paulo. V. 8. N 6. Nov / Dec. 1998.
- MCARDLE, William D., KATCH, Frank I., KATCH, Victor L. Exercise Physiology: Energy, Nutrition and Human Performance. 5th ed. Rio de Janeiro: Guanabara Koogan, 2003.
- MONTEIRO, Wallace. Personal Training: Handbook for Assessment of Prescription Physical Training. 4th ed. Rio de Janeiro: Sprint, 1998.
- OROPEZA, Elsa Silva, REYES, Esperanza Garcia, HERNANDEZ, Lydia Rodriguez; PALENCIA, Luiza Beirana. Mitochondrial Hypertrophic Cardiomyopathy Associated with Wolff-Parkinson-White. Journal of Cardiology. V.82, No. 5, May, 2004.
- RIQUE, Ana Beatriz Ribeiro, SOARES, Eliana de Abreu, MEIRELLES, Claudia de Mello. Nutrition and Exercise in the Prevention of Cardiovascular Disease. Journal of Medicine and Sports. V. 8, No. 6. Nov / Dec, 2002.
- SORBO MD; PLUG GF; Miorelli M, S tered; PERRONE C; LAME S, GRASSO F; GM GIORDANO, A. NAVA The prevalence of Wolff-Parkinson-White Syndrome in a Population of 116,542 young evils. G Ital Cardiol 1995; 25: 681-7.

BRAZILIAN SOCIETY OF hypertension at. Hypertension. Available at: < <http://www.sbh.org.br/geral/hipertensao.asp>> Accessed: 08/07/2011.

VAN Eyken, Braga Elisa Beatriz Del 'Orto. Prevalence of smoking, physical inactivity, Obesity and Hypertension in Adult Men of a Judge of Urban Area Fora, Minas Gerais. Dissertation University Estacio de Sa, 2007.

VARELLA Dr. Dráuzio. Physical activity, diet and diabetes. Published in 00/00/2005, Available at: << <http://www.drauziovarella.com.br/ExibirConteudo/368/atividade-fisica-habitos-alimentares-e-Diabetes>>> Accessed: 07/05/2008.

VILARES, SMF, RIBEIRO, MM, SILVA, AG; PRADO, DML; ZANQUETTA, MM Childhood Obesity. IN: NEGRÃO EC; BARRETO, ACP Cardiology Exercise: From Athlete Heart Disease. 3rd Ed Barueri, SP: Manole, 2010.

ZIMERMAN L.I., FENELON G., MARTINELLI FILHO, M., GRUPI C, LITA J., A. Lorge SON, et al. Brazilian Society of Cardiology. Brazilian Guidelines on Atrial Fibrillation. Arq Bras. Cardiol. 2009, 92 (suppl. 1 6): 1-39

WALLACE, Janet P. Exercise in Hypertension - A Clinical Review. Sports Medicine. V. 33, No. 8, 2003.

WILMORE, Jack H, Costill, David L. Physiology of Sport and Exercise. 1st ed. Manole, Sao Paulo.

FABIO PROCÓPIO SOARES BENTO

Padre Eustáquio Street, 50.

Neighborhood Altino Barbosa

Teófilo Otoni -MG - (33) 3521 0652

COMBINED EFFECT OF TRAINING IN HYPERTENSIVE, OVERWEIGHT, POST-ABLATION SYNDROME WOLFF-PARKINSON-WHITE ABOUT FACTORS THAT TRIGGER ATRIAL FIBRILLATION.

SUMMARY

Exercise training has been recommended as the main agent in the treatment of atrial fibrillation by reducing the weight, control high blood pressure and thus decrease its action. Aims to determine the effects of combined training program an individual with overweight, hypertension, post-ablation of Wolff-Parkinson-White on factors that trigger atrial fibrillation. The training was conducted over a period of 3 months, and the individual has undergone an aerobic and resistance training at moderate intensity. There was a good evolution of the individual with a reduction in BMI from 26.1 to 23.6. Blood pressure was observed during the whole period there has been a slight decrease to levels where significant, the individual did not feel ill during or after exercise. This concludes that the combined training and moderate intensity exercise but significant effect in reducing the weight and in controlling atrial fibrillation.

KEYWORDS: Combined Training, Wolff-Parkinson-White syndrome, atrial fibrillation.

EFFET COMBINE DE FORMATION EN HYPERTENDUS, OBESES, POST-ABLATION SYNDROME DE WOLFF-PARKINSON-WHITE SUR LES FACTEURS QUI DÉCLENCHENT LA FIBRILLATION AURICULAIRE.

SOMMAIRE

L'entraînement physique a été recommandé que l'agent principal dans le traitement de la fibrillation auriculaire en réduisant le poids, le contrôle de l'hypertension artérielle et ainsi diminuer son action. Visé à déterminer les effets du programme de formation combinée d'un individu à l'embonpoint, l'hypertension, post-ablation de Wolff-Parkinson-White sur les facteurs qui déclenchent la fibrillation auriculaire. La formation a été menée sur une période de 3 mois, et l'individu a subi une aérobic et musculation à intensité modérée. Il y avait une bonne évolution de l'individu avec une réduction de l'IMC de 26,1 à 23,6. La pression artérielle a été observée pendant toute la période, il y a eu une légère baisse à des niveaux où significative, l'individu ne se sent pas malade pendant ou après l'exercice. Ceci conclut que la formation combinée et l'exercice d'intensité modérée, mais d'effet significatif dans la réduction du poids et à contrôler la fibrillation auriculaire.

MOTS-CLÉS: Combiné Formation, de Wolff-Parkinson-White Syndrome, une fibrillation auriculaire.

EFFECTO COMBINADO DE FORMACIÓN EN HIPERTENSOS, CON SOBREPESO, DESPUÉS DE LA ABLACIÓN CONEL SÍNDROME DE WOLFF-PARKINSON-WHITE ACERCA DE LOS FACTORES QUE DESENCADENAN LA FIBRILACIÓN AURICULAR.

RESUMEN

La práctica de ejercicio ha sido recomendado como el principal agente en el tratamiento de la fibrilación auricular mediante la reducción del peso, controlar la presión arterial alta por lo tanto disminuir su acción. Tiene como objetivo determinar los efectos del programa de entrenamiento combinado de una persona con sobrepeso, hipertensión, después de la ablación de Wolff-Parkinson-White en los factores que desencadenan la fibrilación auricular. La capacitación se llevó a cabo durante un periodo de 3 meses, y el individuo ha experimentado un entrenamiento aeróbico y de resistencia de intensidad moderada. Hubo una buena evolución de la persona con una reducción en el IMC de 26,1 a 23,6. La presión arterial se observó durante todo el periodo se ha producido un ligero descenso a niveles en los importantes, el individuo no se siente mal durante o después del ejercicio. Esto concluye que el entrenamiento combinado y ejercicio de intensidad moderada, pero un efecto significativo en la reducción del peso y en el control de la fibrilación auricular.

PALABRAS CLAVE: Formación combinada, síndrome de Wolff-Parkinson-White, la fibrilación auricular.

EFEITO DO TREINAMENTO COMBINADO EM INDIVÍDUO HIPERTENSO, COM SOBREPESO, PÓS-ABLAÇÃO DE SÍNDROME DE WOLFF-PARKINSON-WHITE, SOBRE FATORES QUE DESENCADAM A FIBRILAÇÃO ATRIAL.

RESUMO

A prática de exercício tem sido recomendada como principal agente no tratamento da Fibrilação Atrial, por reduzir o peso, controlar a hipertensão arterial e assim diminuir sua ação. Tem como objetivo verificar os efeitos de programa de treinamento combinado um indivíduo com sobrepeso, hipertenso, pós-ablação da Síndrome de Wolff-Parkinson-White, sobre fatores que desencadeiam a Fibrilação Atrial. O treino foi realizado num período de 3 meses, e o indivíduo se submeteu a um treinamento aeróbico e de resistência em intensidade moderada. Verificou-se uma boa evolução do indivíduo com redução do IMC de 26,1 para 23,6. A Pressão Arterial foi observada durante todo o período e notou-se uma leve redução a níveis consideráveis onde, o indivíduo não se sentiu mal, durante ou mesmo depois do exercício. Com isso conclui-se que o treinamento combinado e de intensidade moderada exerce sim efeito significativo na diminuição do peso e no controle da fibrilação atrial.

PALAVRAS-CHAVE: Treinamento Combinado; Síndrome de Wolff-Parkinson-White; Fibrilação Atrial.